

Oral presentation

Open Access

A comprehensive model of idiopathic scoliosis (IS) progression, based on the patho-biomechanics of the deforming "three joint complex"

TB Grivas*¹, ES Vasiliadis², G Triantafyllopoulos¹ and A Kaspiris²

Address: ¹Department of Trauma and Orthopaedics, "Tzanio" General Hospital - NHS, Tzani and Afendouli, 18536, Piraeus, Greece and ²Department of Trauma and Orthopaedics, "Thrasio" General Hospital - NHS, G, Gennimata Av, 19600, Magoula, Attica, Greece D. Bernardou 31 str. Brillissia, 15235 Attica, Greece, 15235, Greece

Email: TB Grivas* - grivastb@vodafone.net.gr

* Corresponding author

from 6th International Conference on Conservative Management of Spinal Deformities
Lyon, France. 21-23 May 2009

Published: 14 December 2009

Scoliosis 2009, 4(Suppl 2):O10 doi:10.1186/1748-7161-4-S2-O10

This abstract is available from: <http://www.scoliosisjournal.com/content/4/S2/O10>

© 2009 Grivas et al; licensee BioMed Central Ltd.

Introduction

It was previously postulated that the intervertebral disc wedging is a significant progressive factor for mild idiopathic scoliosis (IS) curves. The present report introduces an innovative comprehensive model of IS curve progression based on intervertebral disc diurnal variation and the subsequent patho-biomechanics of the deforming "three joint complex".

Methods and materials

Throughout the day and night, due to sustained loading and unloading, the wedged intervertebral disc space in the scoliosis patient expels fluid and reabsorbs it more on the convex side. The convex side of the disc sustains a greater amount of cyclic expansion than the concave side.

Results

Consequently, the imposed convex-wise, asymmetrically concentrated cyclical loads on the adjacent immature vertebral end plates and posterior elements of the spine lead to asymmetrical vertebral growth. More specifically the loading on the two facet-joints asymmetrically increases during the day, as the wedged disc space narrows due to expelled water and it asymmetrically decreases during the night, as the disc space swells due to reabsorbed water.

Discussion

This 24 hour period of cyclic asymmetric loading leads both to asymmetric growth of the end plates and wedging of the vertebral bodies, and to similarly asymmetric growth of the pedicles and arches posteriorly as an effect of Hütter-Volkman law. It is well described that the pedicle in the convex side is more elongated than in the concave side and the facet joint larger respectively.

Significance

The proposed model may help to explain the beneficial effects of exercises, night time bracing in idiopathic scoliosis and of fusionless surgery with staples for progressive IS [1-3].

References

1. Shea KG, Ford T, Bloebaum RD, D'Astous J, King H: **A comparison of the microarchitectural bone adaptations of the concave and convex thoracic spinal facets in idiopathic scoliosis.** *J Bone Joint Surg Am* 2004, **86-A(5)**:1000-6.
2. Grivas TB, Vasiliadis E, Malakasis M, Mouzakis V, Segos D: **Intervertebral disc biomechanics in the pathogenesis of idiopathic scoliosis.** *Stud Health Technol Inform* 2006, **123**:80-3.
3. Iatridis JC, MacLean JJ, O'Brien M, Stokes IA: **Measurements of proteoglycan and water content distribution in human lumbar intervertebral discs.** *Spine* 2007, **32(14)**:1493-7.