

ORAL PRESENTATION

Open Access

# Cheneau bracing with dobomed physiotherapy for thoracic scoliosis: prospective evaluation of 25 patients followed to skeletal maturity

Jacek Durmala\*, Tomasz Kotwicki, J Piotrowski

From 7th International Conference on Conservative Management of Spinal Deformities  
Montreal, Canada. 20-22 May 2010

## Introduction

Giving evidence of efficacy of conservative management of progressive idiopathic scoliosis in the period of rapid adolescent growth requires prospective evaluation and a long time observation, thus the final number of patients available for analysis is often limited.

The aim of the study was to present a series of patients who finished the conservative treatment.

## Material and methods

Twenty-five girls aged 10 to 14 years underwent conservative treatment for thoracic idiopathic scoliosis. All revealed radiological proof of progression. The initial thoracic Cobb angle revealed the value of 16.0° to 40.0°, mean 26.1° ± 8.4°. Cheneau brace was ordered for full time wearing, accompanied with DoboMed daily physiotherapy. The treatment was endorsed during a 2 week in-patient stay at the rehabilitation department; the brace was fitted and the patients learned physiotherapy. Regular controls were performed with a radiological check once a year. The duration of therapy was 36 to 89 months, mean 53.6 ± 14 months. The last radiography was done 6 months after discharging from brace.

## Results

The effective time of brace wearing reported on investigation varied from 8 to 23 hours, mean 11.9 ± 5.5 hours. The Cobb angle at follow-up was 8.0° to 54.0°, mean 32.0° ± 12.7°. The stabilization was achieved in 56% of patients. Three patients (12%) exceeded the value of 50° of Cobb, considered to be surgical indication.

## Conclusion

Stabilization of progressive thoracic scoliosis was achieved in girls using the Cheneau brace and specific DoboMed physiotherapy.

Published: 10 September 2010

doi:10.1186/1748-7161-5-S1-O75

Cite this article as: Durmala et al.: Cheneau bracing with dobomed physiotherapy for thoracic scoliosis: prospective evaluation of 25 patients followed to skeletal maturity. *Scoliosis* 2010 **5**(Suppl 1):O75.

Submit your next manuscript to BioMed Central and take full advantage of:

- Convenient online submission
- Thorough peer review
- No space constraints or color figure charges
- Immediate publication on acceptance
- Inclusion in PubMed, CAS, Scopus and Google Scholar
- Research which is freely available for redistribution

Submit your manuscript at  
www.biomedcentral.com/submit



University of Medical Sciences, Poznan, Poland  
Full list of author information is available at the end of the article