

ORAL PRESENTATION

Open Access

Trunk Appearance Perception Scale (TAPS) discrepancy between scoliosis children and their parents influence the SRS-22 score

M Rigo^{*}, E D'agata, M Jelacic

From 8th International Conference on Conservative Management of Spinal Deformities and SOSORT 2011 Annual Meeting
Barcelona, Spain. 19-21 May 2011

Background

The Trunk Appearance Perception Scale (TAPS) is a valid instrument to assess self perception of trunk deformity [1]. The SRS-22 has been widely used to measure Health Related Quality of Life in scoliosis population but it is not clear which factors can influence its final score [2-4].

Study

Trunk deformity can be perceived differently by children and parents [5,6]. The aim of this study was to check whether different perception of the trunk deformity between patients and parents is a factor influencing the SRS-22 or not.

Materials and methods

Prospective study including 71 (62 F, 9 M) patients with idiopathic scoliosis (treated and non treated), attending the clinic with their parents. Mean age $17 \text{ y} \pm 5.7$. Mean Cobb angle $37^\circ \pm 15^\circ$. All patients completed the SRS-22 and the TAPS. Parents completed the TAPS assessing trunk deformity of their children. A coefficient of discrepancy (TAPS-CD) was defined. Statistical analysis was performed by using SPSS to compare TAPS, TAPS-CD and SRS-22.

Results

Results showed a negative correlation between TPAS-CD and the total SRS-22 ($p < .05$) and treatment satisfaction ($p < .05$). A significant positive correlation was found between patients TAPS, self-image and pain and between parents TAPS, function and treatment satisfaction in the

SRS-22. Two groups were created according to the SRS-22 score. Patients with higher score in the SRS-22 showed a higher TAPS-CD ($P < .05$).

Conclusion

Discrepancy in Trunk Deformity Perception between children and parents influence the SRS-22 of the children. Patients' perception and parents' perception of trunk deformity influence the SRS-22 differently.

Published: 27 January 2012

References

1. Bagó J, Sánchez-Raya J, Pérez-Gruoso FJ, Sánchez, Climent JM: **The Trunk Appearance Perception Scale (TAPS): A new tool to evaluate subjective impression of trunk deformity in patients with idiopathic scoliosis.** *Scoliosis* 2010, **5**:6.
2. Asher M, Lai SM, Burton D, Manna B: **The influence of spinal and trunk deformity on preoperative idiopathic scoliosis patient's health-related quality of life questionnaire responses.** *Spine* 2004, **29**:861-888.
3. Bago J, Climent JM, Ey A, Perez-Gruoso FJ, Izquierdo E: **The Spanish version of the SRS-22 patient questionnaire for idiopathic scoliosis: transcultural adaptation and reliability analysis.** *Spine* 2004, **29**(15):1676-1680.
4. Climent JM, Bagó J, Sánchez J, et al: **Validity of the Spanish version of the Scoliosis Research Society -22 (SRS-22) Patient Questionnaire.** *Spine* 2005, **30**:705-709.
5. Rinella A, Lenke L, Peelle M, Edwards C, Bridwell KH, Sides B: **Comparison of SRS questionnaire results submitted by both parents and patients in the operative treatment of idiopathic scoliosis.** *Spine* 2004, **29**(3):303-310.
6. Smith PL, Donaldson S, Hedden D, Alman B, Howard A, Stephens D, et al: **Parents' and patients' perception of postoperative appearance in adolescent idiopathic scoliosis.** *Spine* 2006, **31**(29):2367-2347.

doi:10.1186/1748-7161-7-S1-O3

Cite this article as: Rigo et al.: Trunk Appearance Perception Scale (TAPS) discrepancy between scoliosis children and their parents influence the SRS-22 score. *Scoliosis* 2012 **7**(Suppl 1):O3.