

Oral presentation

Open Access

Repeatability of the Aesthetic Index for adolescent scoliosis idiopathic evaluation

Fabio Zaina, Stefano Negrini*, Marco Monticone, Chiara Paroli and Angelo Aulisa

Address: ISICO (Italian Scientific Spine Institute), Via Carlo Crivelli 20, 20122 Milan, Italy

Email: Stefano Negrini* - stefano.negrini@isico.it

* Corresponding author

from 4th International Conference on Conservative Management of Spinal Deformities
Boston, MA, USA. 13–16 May 2007

Published: 12 October 2007

Scoliosis 2007, **2**(Suppl 1):S46 doi:10.1186/1748-7161-2-S1-S46

This abstract is available from: <http://www.scoliosisjournal.com/content/2/S1/S46>

© 2007 Zaina et al; licensee BioMed Central Ltd.

Objective

To evaluate the intra- and inter-rater repeatability of the Aesthetic Index (AI) and its sub-scores.

Study design

Since many years in our Institute we have used a clinical tool to evaluate the aesthetics of the trunks posteriorly. This is based on a three point scale for asymmetry (0 absent, 1 slight, 2 important) of the shoulders, scapulae and flanks; the sum of these sub-scores gives the AI. One hundred sixty posterior anterior photographs of the trunk of adolescent idiopathic scoliosis patients were scored two times independently by three observers. We used the Kappa statistics [1]: 0–0.2 poor, 0.2–0.4 fair, 0.4–0.6 moderate, 0.6–0.8 good, 0.8–1.0 very good. The 95% level of agreement was used to identify the minimum clinically significant change to be considered between two different clinical examinations.

Results

See Table 1.

Conclusion

The intra-raters agreement is between moderate and good for single sub-scores, and fair for the AI, as it happens for the inter-raters agreement. The AI and its sub-scores can be used in the everyday clinical practice, but a change of 2 points for sub-scores, and 3 for AI is required to reach clinical significance, and this gives to the AI a low sensitivity.

References

1. Altman DG: **Practical Statistics for Medical Research**. London: Chapman and Hall; 1991.

Table 1: Results.

	Kappa statistics range		Percent of agreement		95% level of agreement (range)	
	Intra-raters	Inter-raters	Intra-raters	Inter-raters	Intra-raters	Inter-raters
Aesthetic Index	0.28–0.41	0.17–0.28	46.2–55.6	37.6–41.5	2(98.7–99.4%)	2 (92.5–99.4%)
Shoulders	0.42–0.53	0.30–0.35	64.4–73.7	58.3–61.2	1 (99.4–100%)	1 (96.9–100%)
Scapulae	0.50–0.58	0.20–0.43	69.4–76.9	53.5–61.9	1 (98.1–100%)	1 (95.6–100%)
Flanks	0.48–0.70	0.28–0.33	75.6–82.5	62.8–66.1	1 (100–100%)	1 (98.7–100%)