

Oral presentation

Open Access

Brace treatment with progressive lordotic forces at the thoracolumbar junction in adolescent scoliosis and hyperkyphosis

Piet JM van Loon*, Monique Roukens and DirkJan Wever

Address: Slingeland Ziekenhuis Doetinchem, Kruisbergseweg 25, Postbus 169, 7000AD, Doetinchem, The Netherlands

* Corresponding author

from 5th International Conference on Conservative Management of Spinal Deformities
Athens, Greece. 3–5 April 2008

Published: 15 January 2009

Scoliosis 2009, **4**(Suppl 1):O47 doi:10.1186/1748-7161-4-S1-O47

This abstract is available from: <http://www.scoliosisjournal.com/content/4/S1/O47>

© 2009 van Loon et al; licensee BioMed Central Ltd.

Introduction

Brace treatment options in idiopathic spinal deformities are limited. Most correctional forces are aimed at the apices of curves in scoliosis and in kyphosis. Psychological drawbacks and discomfort in stiff, unnatural designed orthoses result in unsatisfactory compliance.

Aim

To offer an alternative to present braces. To show that restoration of natural lordosis and concomitant mobility at the thoracolumbar junction by symmetrical lordotic forces, thus only in the sagittal plane, in an adjustable brace offers an interesting alternative.

Study

Retrospective case control study with radiological results and scores for overall-satisfaction.

Methods

Review of 91 children with scoliotic and kyphotic spinal deformities wearing a lordotic brace during growth for at least a full year. Measurements of Cobb angles on AP and sagittal standing X-ray's were compared at indication time, first-in brace day and out of brace after a full year. A questionnaire was filled in with scores for satisfaction, compliance and repeated choice.

Results

Mean age of starting brace was late: 13,8 years (SD2).

In kyphosis (pure or concomitant scoliosis maximum 25°, N = 79)° values for the thoracic curve ($p < 0.01$), the

thoracolumbar curve ($p < 0.01$), de lumbar lordosis ($p < 0.01$) and the pelvic incidence ($p < 0.01$) changed significantly in a paired t-test at one full year brace treatment in comparison with the values at time of indication.

For scoliosis (one curve at least 25°, N = 38) the in-brace correction is significant ($p < 0.01$) of the Cobb angles of thoracic and thoracolumbar curves and the pelvic obliquity. In the sagittal plane even after a full year, a significant correction was seen in the thoracic and thoracolumbar curves. Satisfied and very satisfied were 84.6%. Choice for same treatment was 75.9%.

Conclusion

This study supports an alternative brace technique for treatment of scoliosis and kyphosis with pure lordosis at the thoracolumbar joint, including periodical adjustments combining passive and active components in redressing tensions. Significant reduction of scoliotic and kyphotic curves is possible during growth. By stepwise restoration of thoracolumbar lordosis and preventing overload in compression during sitting creates improved conditions. Good compliance and satisfaction seems part of the process.

References

1. Shelton YA: **Scoliosis and kyphosis in adolescents: diagnosis and management.** *Adolesc Med State Art Rev* 2007, **18**(1):121-39.
2. Negrini S: **Bracing adolescent idiopathic scoliosis today.** *Disabil Rehabil Assist Technol* 2008, **3**(3):107-11.
3. Kotwicki T, Kinel E, Stryla W, Szulc A: **Discrepancy in clinical versus radiological parameters describing deformity due to brace treatment for moderate idiopathic scoliosis.** *Scoliosis* 2007, **2**:18.

4. Roth M: **Idiopathic scoliosis caused by a short spinal cord.** *Acta Radiol Diagn (Stockh)* 1968, **7(3)**:257-71.
5. Roth M: **[Relative osteoneural growth, a theory of osseous deformities].** *Radiol Diagn (Berl)* 1970, **11(6)**:715-24.
6. van Loon PJ, Kuhbauch BA, Thunnissen FB: **Forced lordosis on the thoracolumbar junction can correct coronal plane deformity in adolescents with double major curve pattern idiopathic scoliosis.** *Spine* 2008, **33(7)**:797-801.

Publish with **BioMed Central** and every scientist can read your work free of charge

"BioMed Central will be the most significant development for disseminating the results of biomedical research in our lifetime."

Sir Paul Nurse, Cancer Research UK

Your research papers will be:

- available free of charge to the entire biomedical community
- peer reviewed and published immediately upon acceptance
- cited in PubMed and archived on PubMed Central
- yours — you keep the copyright

Submit your manuscript here:
http://www.biomedcentral.com/info/publishing_adv.asp

