

ORAL PRESENTATION

Open Access

Early results of Rigo-Chêneau type brace treatment

T Maruyama*, Y Nakao, H Yamada

From 8th International Conference on Conservative Management of Spinal Deformities and SOSORT 2011 Annual Meeting
Barcelona, Spain. 19-21 May 2011

Background

We have been using Rigo-Chêneau type brace for the treatment of idiopathic scoliosis since 2007 [1,2]. Curves other than the upper thoracic main curve were the subjects of the treatment. Most patients wore their brace as part-time, at home or at night.

Purpose

To evaluate early results of Rigo-Chêneau type brace treatment.

Materials and methods

A total of 54 patients, 49 females and 5 males, were included in the analysis. Average age at the beginning of the treatment was 12.5 years (10 to 15). Risser sign was 0 in 15, I in 9, II in 15, III in 5 and IV in 10 patients. Curve pattern was thoracic (T) in 25, thoracolumbar or lumbar (TL) in 10 and double (D) in 19 patients.

Results

Average Cobb angle before treatment was 36.5°, which was reduced in the trial brace to 23.2°: correction rate was 36% (34% for T, 69% for TL, and 31% for D curve). Of 54 patients, 20 met the inclusion criteria of the SRS brace study (Risser 0-2, Cobb angle 25-40°) and six of them reached skeletal maturity during the treatment period. Three of them (50%) progressed more than 6°; however, only one patient progressed more than 10°.

Conclusions

Although early experience suggested better results than the natural history, accumulation of data will be

necessary to determine the effectiveness of the treatment with Rigo-Chêneau type brace.

Published: 27 January 2012

References

1. Rigo M, Weiss HR: The Cheneau concept of bracing- Biomechanical aspects. *Stud Health Technol and Inform* 2008, **135**:303-319.
2. Rigo M, Villagrasa M, Gallo D: A specific scoliosis classification correlating with brace treatment: description and reliability. *Scoliosis* 2010, **5**:1.

doi:10.1186/1748-7161-7-S1-O33

Cite this article as: Maruyama et al.: Early results of Rigo-Chêneau type brace treatment. *Scoliosis* 2012 **7**(Suppl 1):O33.

Submit your next manuscript to BioMed Central and take full advantage of:

- Convenient online submission
- Thorough peer review
- No space constraints or color figure charges
- Immediate publication on acceptance
- Inclusion in PubMed, CAS, Scopus and Google Scholar
- Research which is freely available for redistribution

Submit your manuscript at
www.biomedcentral.com/submit



Saitama Medical Center, Saitama Medical University, Kawagoe, Japan