

ORAL PRESENTATION

Open Access

Scoliosis and sagittal balance in Parkinson's disease: analysis of correlations

Luciano Bissolotti*, Massimiliano Gobbo, Fabio Zaina, Monia Lusini, Sabina Donzelli, Stefano Negrini

From 10th International Conference on Conservative Management of Spinal Deformities - SOSORT 2013 Annual Meeting

Chicago, IL, USA. 8-11 May 2013

Background

Information concerning scoliosis in Parkinson's disease (PD) and its correlations with sagittal balance (SB) is sparse.

Purpose

The aim of this study was to describe the prevalence of scoliosis in PD patients and the existing correlations with SB in relation to the spinopelvic morphology.

Methods

A total of 48 consecutive PD patients were included: 36 males, 12 females; 70.8±7.6 years; 6.4±4.1 years of disease (YOD); Hoehn Yahr (HY) 2.7±1.2. The clinical assessment included HY score, Pain NRS 0-10 and trunk rotation in bending (ATR). Lumbar lordosis (LL), thoracic kyphosis (TK), scoliosis curves (SC), spinosacral angle (SSA), spinopelvic angle (SPA), pelvic incidence (PI), sacral slope (SS) and pelvic tilt (PT) were radiographically assessed. Patients have been compared according to the presence of SC >10° (PD_{ts}) Cobb or the absence of SC (PD_{ns}).

Results

Among the study subjects, 47.9% presented a SC larger than 10°, 84% of the patients in PD_{ts} presented a thoracolumbar curve, 10% a thoracic curve and 6% a lumbar curve. The cohort did not present differences with PD_{ns} about age (71.8±6.0 vs. 69.8±8.8yrs) and YOD (6.1±4.1 vs. 6.6±4.1 years). No differences have been detected for HY score (2.7±1.2 vs. 2.6±1.6) and NRS (29.6±22.6 vs. 19.4±28.1). ATR was higher in PD_{ts} (5.6±4.9 vs. 1.3±1.9, p<0.01). TK (46.4±16.1 vs. 46.9±12.1°), LL (46.3±26.9 vs. 49.3±13.9°), SSA (104.8±24.7 vs. 118.6±12.9°) and SPA

(152.4±20.3 vs. 153.4±12.5°) were not different (p>0.05). PI (57.8±11.1 vs. 53.9±13.1°) and PT (23.6±13.7 vs. 17.6±8.6°) were slightly but not statistically different, while SS was not (35.3±12.1 vs. 36.0±8.5°).

Conclusions and discussion

The prevalence of scoliosis in PD was higher than previously described by other authors, with the thoracolumbar spine mostly affected. SB was not different between two groups while, in PD_{ts}, spinopelvic parameters presented the tendency to have a larger PI and PT.

Published: 18 September 2013

References

1. Koller H, Acosta F, Zenner J, Ferraris L, Hitzl W, Meier O, Ondra S, Koski T, Schmidt R: Spinal surgery in patients with Parkinson's disease: experiences with the challenges posed by sagittal imbalance and the Parkinson's spine. *Spine J* 2010, **19**(10):1785-94, Epub 2010 Apr 27.
2. Doherty KM, van de Warrenburg BP, Peralta MC, Silveira-Moriyama L, Azulay JP, Gershanik OS, Bloem BR: Postural deformities in Parkinson's disease. *Lancet Neurol* 2011, **10**(6):538-549.
3. Baik JS, Kim JY, Park JH, Han SW, Park JH, Lee MS: Scoliosis in patients with Parkinson's disease. *J Clin Neurol* 2009, **5**(2):91-94.

doi:10.1186/1748-7161-8-S2-O7

Cite this article as: Bissolotti et al.: Scoliosis and sagittal balance in Parkinson's disease: analysis of correlations. *Scoliosis* 2013 **8**(Suppl 2):O7.

* Correspondence: Luciano.bissolotti@ancelle.it
Rehabilitation Service, Casa di Cura Domus Salutis, Brescia, Italy